

Anti-DENV NS4B Antibody [Clone 24-1]

Catalog Number LDG0008YA

Package 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Overview

Description

Dengue virus belongs to the family Flaviviridae, genus Flavivirus and consists of four distinct serotypes (DENV1 to 4). Dengue virus is transmitted by mosquitos majorly of species *Aedes aegypti* and *Aedes Albopictus* that widespread around tropics and subtropics. NS4B is a non-enzymatic transmembrane protein, acting as an important component of the viral replication complex in the ER. The NS4B proteins can affect host immunity through protein-protein interactions.

Product Note

Recommended dilution factor:

ELISA: 1:5000-20000

WB: 1:1000-5000

IFA: 1:500-1000

FACS: Assay dependent

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Specifications

Host

Mouse

Isotype

IgG1

Clonality

Monoclonal

Clone Name

clone 24-1

Immunogen

Dengue virus NS4B

Application

ELISA, WB, IFA, FACS

Concentration

1 mg/mL

Specificity

NS4B

Reactivity

Dengue virus

Conjugation

Unconjugated

Storage Buffer

Phosphate Buffered Saline containing 0.03% ProClin 300, pH 7.4.

Form

Liquid

Instruction**Shipping**

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

This product is stable after storage at:


- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles.


Suggestion: Divide antibody into several vials.

Keep only vials for usage at 2-8°C.


Image




Immunofluorescence analysis of Anti-DENV NS4B Antibody [Clone 24-1]
Control BHK cells (uninfected) and Virus (DENV 2) infection BHK cells were fixed in 4% PFA, permeabilized with PBS containing 0.1% Triton X-100. Cells were stained with mouse anti-DENV NS4B monoclonal antibody (1:400) followed by secondary antibodies (goat anti-Mouse IgG-iFluor 488, 1:400, green) and cell nuclei were stained with Hoechst 33342 (Blue).



Immunofluorescence analysis of Anti-DENV NS4B Antibody [Clone 24-1]
Control Vero cells (uninfected) and Virus (DENV 2,3) infection Vero cells were fixed in 4% PFA, permeabilized with PBS containing 0.1% Triton X-100. Cells were stained with mouse anti-DENV NS4B monoclonal antibody (1:400) followed by secondary antibodies (goat anti-Mouse IgG-iFluor 488, 1:400, green) and cell nuclei were stained with Hoechst 33342 (Blue).



ELISA titration of Anti-DENV NS4B Antibody [Clone 24-1]
Titration curve of anti-DENV NS4B antibody in ELISA. Red: NS4B; Black: BSA (negative control).



Western blotting analysis of Anti-DENV NS4B Antibody [Clone 24-1]

Western blot: Virus infection BHK cell lysates were stained with mouse anti-DENV NS4B monoclonal antibody at 1:1000 dilution.

Lane 1: BHK cell lysate (10 µg),
Lane 2: ZIKV infection (10 µg),
Lane 3: DENV1 infection (10 µg),
Lane 4: DENV2 infection (10 µg),
Lane 5: DENV3 infection (10 µg),
Lane 6: DENV4 infection (10 µg).

Disclaimer : For Research Use or Further Manufacturing Only.