

## Goat Anti-Human IgG (Fc Fragment) Antibody (HRP)

**Catalog Number** LDG0017YE

**Package** 1 mL / Customized package

For full product information, images and publications, please visit [our website](#).



### Overview

#### Description

Leadgene Goat anti-Human IgG, Fc Fragment is purified from goat antiserum by immunoaffinity chromatography and conjugated with high-activity HRP. Minimal cross-reactivity with other isotypes or species.

#### Product Note

Recommended dilution factor:

ELISA: 1:5000-25000

IHC: 1:500-2500

WB: 1:5000-25000

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

### Specifications

#### Host

Goat

#### Immunogen

Gamma immunoglobulin Fc region

#### Application

ELISA, IHC, WB

#### Clonality

Polyclonal

#### Reactivity

Human

#### Conjugation

HRP

**Storage Buffer**

Phosphate Buffered Saline containing 1% BSA, 0.03% ProClin 300 and 10% glycerol, pH 7.4.

**Form**

Liquid

**Specificity**

IgG, Fc Fragment

**Instruction****Shipping**

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

**Stability & Storage**

This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles and protect from light.

Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

**Disclaimer :** For Research Use or Further Manufacturing Only.