

Murine RNase Inhibitor

Catalog Number	LDG0018RI
Package	4 KU / 20 KU / Customized package

For full product information, images and publications, please visit [our website](#).



Overview

Description

Murine RNase Inhibitor effectively inhibits various RNases (RNase A, B, C). Its superior antioxidant activity, due to the absence of two oxidation-sensitive cysteine residues found in human inhibitors, makes it ideal for DTT-sensitive experiments like qPCR. Additionally, it is free from DNase, RNase, and Nickase contaminants, ensuring RNA sample integrity.

Specifications

Expression System

Escherichia coli

Storage Buffer

20 mM HEPES-KOH, 50 mM KCl, 8 mM DTT, 50% glycerol, pH 7.6

Unit Definition

One unit is defined as the amount of Murine RNase Inhibitor required to inhibit the activity of 5 ng of RNase A by 50%. Activity is measured by the inhibition of hydrolysis of cytidine 2', 3'-cyclic monophosphate by RNase A.

Concentration

40 U/ μ L

Purity

>95% as determined by SDS-PAGE analysis.

Form

Liquid

Instruction

Tainan Headquarter

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles.

Suggestion: Divide antibody into several vials.

Keep only vials for usage at 2-8°C.
