

## Murine RNase Inhibitor

<b>Catalog Number</b>	LDG0018RI
<b>Package</b>	4 KU / 20 KU / Customized package

For full product information, images and publications, please visit [our website](#).



### Overview

#### Description

Murine RNase Inhibitor effectively inhibits various RNases (RNase A, B, C). Its superior antioxidant activity, due to the absence of two oxidation-sensitive cysteine residues found in human inhibitors, makes it ideal for DTT-sensitive experiments like qPCR. Additionally, it is free from DNase, RNase, and Nickase contaminants, ensuring RNA sample integrity.

### Specifications

#### Expression System

Escherichia coli

#### Concentration

40 U/μL

#### Storage Buffer

20 mM HEPES-KOH, 50 mM KCl, 8 mM DTT, 50% glycerol, pH 7.6

#### Purity

>95% as determined by SDS-PAGE analysis.

#### Unit Definition

One unit is defined as the amount of Murine RNase Inhibitor required to inhibit the activity of 5 ng of RNase A by 50%. Activity is measured by the inhibition of hydrolysis of cytidine 2', 3'-cyclic monophosphate by RNase A.

#### Form

Liquid

### Instruction

#### Tainan Headquarters

+886-6-2536677

bd@leadgene.com.tw

#### Innovation & Research Center

+886-2-27065528

#### CLD Center

+886-6-2536677

---

### Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

---

### Stability & Storage

This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles.

Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

---

---

**Disclaimer :** For Research Use or Further Manufacturing Only.

---

#### Tainan Headquarters

☎ +886-6-2536677

✉ [bd@leadgene.com.tw](mailto:bd@leadgene.com.tw)

---

#### Innovation & Research Center

☎ +886-2-27065528

---

#### CLD Center

☎ +886-6-2536677