

Anti-IL-8 Antibody [Clone 58-1]

Catalog Number LDG0030YA

Package 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Overview

Description

The CXC chemokine is a group of secreted proteins, attracts neutrophils and lymphocytes. Interleukin-8 (IL-8, or CXCL8), a member of the CXC chemokines, is a proinflammatory cytokine produced by macrophages and other cell types like endothelial cells, epithelial cells and airways smooth muscle cells. This cytokine mainly attracts and activates neutrophils to the site of inflammation and also plays an important role in cell proliferation, tissue remodeling and angiogenesis.

Product Note

Recommended dilution factor:

ELISA: 1:5000-20000

WB: 1:1000-5000

IFA: 1:500-1000

FACS: Assay dependent

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Specifications

Host

Mouse

Isotype

IgG1

Clonality

Monoclonal

Clone Name

clone 58-1

Immunogen

Interleukin 8

Application

ELISA, WB, IFA, FACS

Concentration

1 mg/mL

Specificity

Interleukin 8

Reactivity

Human

Conjugation

Unconjugated

Storage Buffer

Phosphate Buffered Saline containing 0.03% ProClin 300, pH 7.4.

Form

Liquid

Instruction**Shipping**

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

This product is stable after storage at:


- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles.


Suggestion: Divide antibody into several vials.

Keep only vials for usage at 2-8°C.


Image




Immunofluorescence analysis of Anti-IL-8 Antibody [Clone 58-1] HeLa cells were fixed in 4% PFA, permeabilized with PBS containing 0.1% Triton X-100. Cells were stained with mouse anti-IL-8 monoclonal antibody (1:200) followed by secondary antibodies (goat anti-Mouse IgG-iFluor 488, 1:400, green) and cell nuclei were stained with Hoechst 33342 (Blue).



FACS analysis of Anti-IL-8 Antibody [Clone 58-1] Human peripheral blood mononuclear cells (PBMC) were stained with mouse anti-IL-8 monoclonal antibody at 2 $\mu\text{g/ml}$ (red) and without antibody control (black).



ELISA titration of Anti-IL-8 Antibody [Clone 58-1] Titration curve of anti-IL-8 antibody in ELISA. Red: IL-8; Black: BSA (negative control).



Western blotting analysis of Anti-IL-8 Antibody [Clone 58-1] Recombinant IL-8 protein (100 ng) was stained with mouse anti-IL-8 monoclonal antibody at 1:5000 dilution.

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677