

Anti-D-Dimer Antibody [Clone DD03]

Catalog Number	LDG0165YA
Package	100 μg / Customized package

For full product information, images and publications, please visit our website.



Overview

Description

Blood clotting is an essential physiological process preventing losing blood and pathogen infection in wounds. Subsequent fibrinolytic lysis, d-dimer is the most miniature fibrinolysis-specific degradation product in circulation. The d-dimer is served as a clinically crucial diagnostic marker for blood clotting disorders such as deep vein thrombosis (DVT), pulmonary embolism (PE), and disseminated intravascular coagulation (DIC). These systemic illnesses can be severe and even life-threatening.

Product Note

Recognize human D-Dimer in ELISA, when monoclonal antibody was paired with Anti-D-Dimer Antibody [clone DD01] (cat. LDG0166YA) or [clone DD02] (cat. LDG0167YA).

Recommended dilution factor:

ELISA: 1:5000-20000

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Specifications		
Host	Clonality	
Mouse	Monoclonal	
Isotype	Clone Name	
IgG2b	clone DD03	

Tainan Headquarter

Innovation & Research Center

CLD Center



Application

ELISA, CLIA, LFIA

Concentration

1 mg/mL

Specificity

D-Dimer

Conjugation

Unconjugated

Storage Buffer

Phosphate Buffered Saline containing 0.03% ProClin 300, pH 7.4.

Form

Liquid

Instruction

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage


This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles. Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

Image





ELISA titration of Anti-D-Dimer Antibody [Clone DD03] The D-Dimer and BSA were coated onto wells of a 96 well plate. Indirect ELISA was performed using Anti-D-Dimer Antibody [Clone DD03].

Disclaimer: For Research Use or Further Manufacturing Only.