

Human FGF-10, His Tag, E. coli

Catalog Number LDG076PHE

 $5~\mu g$ / $20~\mu g$ / $100~\mu g$ / Customized package

For full product information, images and publications, please visit our website.



Specifications

Species of Origin

Human

Package

Affinity Tag

His Tag (C-term)

Purity

>98% as determined by SDS-PAGE analysis.

Activity

Measure by its ability to induce 3T3 cells proliferation. The ED $_{50}$ for this effect is <8 ng/mL. The specific activity of recombinant human FGF-10 is > 1.2 x 10 5 IU/mg.

Form

Lyophilized

Expression System

Escherichia coli

Storage Buffer

Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

Molecular weight

The protein has a calculated MW of 20.06 kDa. The protein migrates as 19 kDa under reducing condition (SDS-PAGE analysis).

Endotoxin Level

 $<\!0.1$ EU per 1 μg of the protein by the LAL method.

Background

Tainan Headquarter

Innovation & Research Center

CLD Center



Background

Fibroblast Growth Factors-10 (FGF-10) is a 23.4 kDa member of the fibroblast Growth Factors with 208 amino acid residues. FGF-10 is mainly secreted from endometrial stromal cells, fibroblasts, peritubular cells, leydig cells, muller glia cells. FGF-10 a multifunctional mesenchymal-epithelial signaling Growth Factors, can regulate embryonic development, cell proliferation and cell differentiation. It has been associated with cancer and human genetic disorders.

Synonyms

Keratinocyte Growth Factors 2, Fibroblast Growth Factors 10

Uniprot ID

#015520

Sequence Note

Leu40-Ser208

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration not less than 200 μ g/mL and incubate the stock solution for at least 20 min to ensure sufficient redissolved.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.


Image

Tainan Headquarter

Innovation & Research Center

CLD Center





SDS-PAGE analysis of recombinant human FGF-10.

Disclaimer: For Research Use or Further Manufacturing Only.