

# Human BMP-2, His Tag, E. coli

Catalog Number LDG099PHE

5 μg / 20 μg / 100 μg / Customized package

For full product information, images and publications, please visit our website.



# **Specifications**

### **Species of Origin**

Human

**Package** 

## **Affinity Tag**

His Tag (C-term)

#### **Purity**

>95% as determined by SDS-PAGE analysis.

### **Activity**

Measure by its ability to induce alkaline phosphatase production by ATDC5 cells. The ED $_{50}$  for this effect is <1.0  $\mu$ g/mL.

**Form** 

Lyophilized

## **Expression System**

Escherichia coli

### **Storage Buffer**

Lyophilized from a 0.2  $\mu m$  filtered solution containing 20 mM sodium citrate and 0.2 M NaCl, pH 4.5.

### Molecular weight

The protein has a calculated MW of 13.76 kDa. The protein migrates as 14 kDa under reducing condition (SDS-PAGE analysis).

### **Endotoxin Level**

<0.1 EU per 1  $\mu$ g of the protein by the LAL method.

# **Background**



## **Background**

Bone Morphogenetic Protein-2 (BMP-2) is an extracellular multifunctional signaling cytokine that is also a member of the TGF- $\beta$  family. BMP-2 can bind with the TGF- $\beta$  receptor to active SMAD protein signal transduction. Moreover, it engages in the path of signals such as Wnt and Notch. BMP-2 plays an essential role in the growth of hard bones and cartilage and involved in the differentiation of osteoblasts.

## **Synonyms**

Bone morphogenetic protein 2A , BMP-2A, Bone morphogenetic protein 2

### **Uniprot ID**

#P12643

### **Sequence Note**

Gln283-Arg396

## Instruction

#### Reconstitution

It is recommended to reconstitute the lyophilized protein in 4 mM HCl to a concentration not less than 200  $\mu$ g/mL and incubate the stock solution for at least 20 min to ensure sufficient redissolved.

# Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

## **Stability & Storage**


This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

## **Image**





SDS-PAGE analysis of recombinant human BMP-2.

**Disclaimer:** For Research Use or Further Manufacturing Only.