

Human FGF-2, Tag free, E. coli

Catalog Number LDG189PHE

Package 5 µg / 20 µg / 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Specifications

Species of Origin

Human

Affinity Tag

Tag Free

Purity

>98% as determined by SDS-PAGE analysis.

Activity

Measure by its ability to induce 3T3 cells proliferation. The ED₅₀ for this effect is <1 ng/mL. The specific activity of recombinant human FGF-2 is approximately >5 x 10⁵ IU/mg.

Form

Lyophilized

Expression System

Escherichia coli

Storage Buffer

Lyophilized from a 0.2 µm filtered solution contain 20 mM Tris, 1 M NaCl, pH 7.4.

Molecular weight

The protein has a calculated MW of 18.1 kDa. The protein migrates as 15-18 kDa under reducing condition (SDS-PAGE analysis).

Endotoxin Level

<0.1 EU per 1 µg of the protein by the LAL method.

Background

Tainan Headquarter

+886-6-2536677

bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528

CLD Center

+886-6-2536677

Background

Basic fibroblast Growth Factors (FGF-2, bFGF, FGF- β), a 18 kDa pleiotropic cytokine, plays multiple roles in different cells and tissues. FGF-2 can stimulate smooth muscle cell growth, wound healing, and tissue repair. In addition, FGF-2 has been shown to regulate the generation of neurons and astrocytes from progenitor cells. FGF-2 are also involved in a variety of biological processes, including embryonic development, morphogenesis, tissue repair, tumor growth, and invasion. As a multifunctional cytokine, FGF-2 is first isolated from the pituitary. Later, it was identified from various cell types including cardiac myocytes, cardiac fibroblasts, endothelial cells, and smooth muscle cells.

Uniprot ID

#P09038

Synonyms

Fibroblast Growth Factors 2, Basic fibroblast Growth Factors , bFGF, Heparin-binding Growth Factors 2 , HBGF-2

Sequence Note

Ala144-Ser288

Instruction

Reconstitution

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 200 μ g/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.


Stability & Storage

This product is stable after storage at:

- -20°C for 12 months in lyophilized state from date of receipt.
- -20°C or -80°C for 1 month under sterile conditions after reconstitution.

Avoid repeated freeze/thaw cycles.

Image



SDS-PAGE analysis of recombinant human FGF-2.

Disclaimer : For Research Use or Further Manufacturing Only.