



# Product Information & Manual

Information of other products is available at: [www.leadgenebio.com](http://www.leadgenebio.com)

## HyFect<sup>®</sup> Transfection Reagent

Cat no. LDG0001RA

### Product Overview

#### Package component

Item	Content
HyFect <sup>®</sup> Transfection Reagent	500 µL

### Description

HyFect<sup>®</sup> Transfection Reagent is a low cell toxicity and non-liposome reagent optimized for DNA delivery. It provides effectively, reproducibly, and affordable benefits for scientific research.

### Storage and Stability

- (1) Stored at -20°C.
- (2) All reagents are stable for one year under proper storage conditions.

### Procedure

#### Standard RNA synthesis procedures:

- (1) Cell preparation: Cells should be seeded before 16 to 20 hours prior to transfection with around 70% confluence. The medium should be refreshed 30 minutes before transfection. Usually, culture medium with serum does not affect transfection.
- (2) DNA preparation: DNA plasmid for transfection should be with high purity (A260/A280=1.8-1.9) to ensure efficient transfection mixture preparation.
- (3) Mixture preparation: Guideline of DNA plasmid and HyFect<sup>®</sup> Transfection Reagent amount and ratio refer to table 1. In brief, plasmid and transfection reagent are diluted in serum-free culture medium for 5 minutes

then mixed together and tip gently for another 25 minutes.

- (4) Transfection: Add mixtures into cell culture dish/plate. The mixture could be removed after 6 to 48 hours and refilled with culture medium.

Recommended formula of transfection mixture

Culture Dish/Plate	Medium Volume	Plasmid / Serum-free medium	HyFect <sup>®</sup> Transfection Reagent / Serum-free medium
96-well	100 µL	250 ng / 10 µL	0.75 µL / 10 µL
24-well	500 µL	500 ng / 25 µL	1.5 µL / 25 µL
12-well	700 µL	750 ng / 35 µL	2.25 µL / 35µL
6-well	1 mL	1 µg / 50 µL	3 µL / 50 µL
6 cm	3 mL	2.5 µg / 150 µL	7.5 µL / 150 µL
10 cm	6 mL	5 µg / 300 µL	15 µL / 300 µL

### Important notes

- (1) HyFect<sup>®</sup> Transfection Reagent is recommended for use with Gibco™ Opti-MEM™ I Reduced-Serum Medium or Gibco™ OptiPRO™ SFM for alternative optimizing transfection efficiency.

### Disclaimer

This product is for research use only and is not intended for diagnostic use

## LEADGENE BIOMEDICAL, INC.

No.9, Ln. 147, Zhengbei 1st Rd., Yongkang Dist., Tainan City 710, Taiwan R.O.C. TEL: +886-6-2536677 FAX: +886-6-2531536

[www.leadgenebio.com](http://www.leadgenebio.com)