

Anti-His Tag Antibody [Clone 14-4]

Catalog Number LDG0001YB

Package 100 μg / Customized package

😳 Publications (3)

Overview

Description

Mouse anti-His tag mAb is specific to 6X His tag, which can be applied to several immunoassays, including ELISA, WB, IP, IFA, and FACS. This antibody has been identified in several sources (bacteria, yeast, and mammalian cells) of His tag fusion protein.

Product Note

Recommended dilution factor: ELISA: 1:5000-20000 WB: 1:1000-5000 IP: 1:500-2000 IFA: 1:500-1000 FACS: 1:500-1000

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Specifications			
Host	Clonality		
Mouse	Monoclonal		
Isotype	Clone Name		
lgG1	clone 14-4		
1901			
Immunogen	Application		
6X His tag fusion protein	ELISA, WB, IP, IFA, FACS		
Tainan Headquarter Innovation & Research	Center CLD Center		
® +886-6-2536677 ® +886-2-27065528	® +886-6-2536677		
☑ bd@leadgene.com.tw			

For full product information, images and publications, please visit our website.





Conjugation

Unconjugated

Buffer

Phosphate Buffered Saline containing 0.03% ProClin 300, pH 7.4.

Form

Liquid

Instruction

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Concentration

1 mg/mL

Specificity

Recognizes 6X His tag on either the N- or Cterminus of recombinant fusion protein.

Stability & Storage

This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles. Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

Image

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw



kDa	1	2	
75-			
63—		-	
48—			
35—			
25—			
17-			

Western blotting analysis of anti-His mAb (1:5000) Lane 1: 293T whole cell lysateuntransfected Lane 2: 293T whole cell lysatetransfected Lysate/proteins at 20 ug per lane

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

Innovation & Research Center

CLD Center

® +886-6-2536677

& +886-2-27065528

& +886-6-2536677

☑ bd@leadgene.com.tw