

Anti-cMyc Tag Antibody [Clone 66-38]

Catalog Number LDG0

Package

LDG0004YB

100 µg / Customized package

For full product information, images and publications, please visit our website.



Overview

Description

Mouse anti-cMyc tag mAb is specific to EQKLISEEDL tags which can be applied to several immunoassays including ELISA, WB, IP, IFA and FACS. This antibody has been identified in several sources (bacteria, yeast, and mammalian cells) of cMyc fusion protein.

Product Note

Recommended dilution factor: ELISA: 1:5000-20000 WB: 1:1000-5000 IP: 1:500-2000 IFA: 1:500-1000 FACS: 1:500-1000

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Specifications	
Host	Clonality
Mouse	Monoclonal
Isotype	Clone Name
lgG1	clone 66-38
Immunogen	Application
Peptide: EQKLISEEDL (cMyc)	ELISA, WB, IP, IFA, FACS

Tainan Headquarter	Innovation & Research Center	CLD Center
	& +886-2-27065528	® +886-6-2536677
☑ bd@leadgene.com.tw		



Conjugation

Unconjugated

Buffer

Phosphate Buffered Saline containing 0.03% ProClin 300, pH 7.4.

Form

Liquid

Concentration

1 mg/mL

Specificity

Recognizes cMyc tag on either the N- or C-terminus of recombinant fusion protein.

Instruction

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles. Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

Image

Tainan Headquarter

Innovation & Research Center

CLD Center

<u>®</u> +886-6-2536677

& +886-2-2706<u>5</u>528

& +886-6-2536677

☑ bd@leadgene.com.tw



75 - 63-	2 TEINIE		
48- 35-	e	LE ADGI	
25			() LEAD

Western blotting analysis of anti-c-Myc tag mAb (1:5000) Lane 1: 293T whole cell lysateuntransfected Lane 2: 293T whole cell lysatetransfected Lysate/proteins at 30 µg per lane

Disclaimer : For Research Use or Further Manufacturing Only.

Tainan Headquarter

Innovation & Research Center

CLD Center

& +886-6-2536677

& +886-6-2536677

☑ bd@leadgene.com.tw