

Anti-DENV NS5 Antibody [Clone 37-2]

Catalog Number LDG0007YA

Package 100 µg / Customized package

For full product information, images and publications, please visit [our website](#).



Overview

Description

Dengue virus belongs to the family Flaviviridae, genus Flavivirus and consists of four distinct serotypes (DENV1 to 4). Dengue virus is transmitted by mosquitos majorly of species Aedes aegypti and Aedes Albopictus that widespread around tropics and subtropics. NS5 is the largest and the most conserved protein of Dengue virus, acting as both a RNA-dependent RNA polymerase and a RNA methyltransferase enzyme. The NS5 proteins are important for viral replication and can down-regulate the host immune interferon response.

Product Note

Recommended dilution factor:

ELISA: 1:5000-20000

WB: 1:1000-5000

IFA: 1:500-1000

FACS: Assay dependent

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Specifications

Host

Mouse

Clonality

Monoclonal

Isotype

IgG2b

Clone Name

clone 37-2

Immunogen

Dengue virus NS5

Application

ELISA, WB, IFA, FACS

Concentration

1 mg/mL

Specificity

NS5

Reactivity

Dengue virus

Conjugation

Unconjugated

Buffer

Phosphate Buffered Saline containing 0.03% ProClin 300, pH 7.4.

Form

Liquid

Instruction**Shipping**

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles.

Suggestion: Divide antibody into several vials.

Keep only vials for usage at 2-8°C.

Image**Tainan Headquarter**

+886-6-2536677


bd@leadgene.com.tw

Innovation & Research Center

+886-2-27065528


CLD Center

+886-6-2536677




Immunofluorescence analysis of Anti-DENV NS5 Antibody [Clone 37-2]

Control Vero cells (uninfected) and Virus (DENV 1,2,4) infection Vero cells were fixed in 4% PFA, permeabilized with PBS containing 0.1% Triton X-100. Cells were stained with mouse anti-DENV NS5 monoclonal antibody followed by secondary antibodies (goat anti-Mouse IgG-iFluor 488, 1:400, green) and cell nuclei were stained with Hoechst 33342 (Blue).



ELISA titration of Anti-DENV NS5 Antibody [Clone 37-2]
 Titration curve of anti-DENV NS5 antibody in ELISA. Red: NS5; Black: BSA (negative control).



Western blotting analysis of Anti-DENV NS5 Antibody [Clone 37-2]
 Virus infection Vero cell lysates were stained with mouse anti-DENV NS5 monoclonal antibody at 1:1000 dilution. Lane 1: Vero cell lysate (10 µg), Lane 2: ZIKV infection (10 µg), Lane 3: DENV1 infection (10 µg), Lane 4: DENV2 infection (10 µg), Lane 5: DENV3 infection (10 µg), Lane 6: DENV4 infection (10 µg).

Disclaimer : For Research Use or Further Manufacturing Only.