

Goat Anti-Human IgG (Fc Fragment) Antibody (HRP)

Catalog Number LDG0017YE

Package 1 mL / Customized package

For full product information, images and publications, please visit our website.



Overview

Description

Leadgene Goat anti-Human IgG, Fc Fragment is purified from goat antiserum by immunoaffinity chromatography and conjugated with high-activity HRP. Minimal cross-reactivity with other isotypes or species.

Product Note

Recommended dilution factor:

ELISA: 1:5000-25000 IHC: 1:500-2500 WB: 1:5000-25000

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

Specifications

Host
Goat
Polyclonal

Immunogen
Reactivity
Gamma immunoglobulin Fc region
Human

Application
ELISA, IHC, WB
HRP

Tainan Headquarter

Innovation & Research Center

CLD Center



Buffer

Phosphate Buffered Saline containing 1% BSA, 0.03% ProClin 300 and 10% glycerol, pH 7.4.

Form

Liquid

Specificity

IgG, Fc Fragment

Instruction

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

This product is stable after storage at:

- 2-8°C for 2 weeks under sterile conditions from date of receipt.
- -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles and protect from light.

Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

Disclaimer: For Research Use or Further Manufacturing Only.