

Bst DNA Polymerase

Catalog Number	LDG0021RF
Package	1,600 U / 8,000 U / Customized package

For full product information, images and publications, please visit our website.



Overview

Description

Bst DNA Polymerase (Large fragment) is an enzyme of Bacillus stearothermophilus DNA polymerase which can catalyze $5' \rightarrow 3'$ polymerase activity but lacks $5' \rightarrow 3'$ exonuclease activity. Bst DNA Polymerase offers strand displacement capabilities, making it ideal for isothermal amplification.

Product Note

- It is not recommended to perform reaction above 70 °C. Bst DNA Polymerase cannot be used for thermal cycle sequencing.
- 10X Bst DNA Polymerase Reaction Buffer: 200 mM Tris-HCl (pH 8.8), 10 mM (NH₄)₂SO₄, 10 mM KCl, 20 mM MgSO₄, and 1% Triton® X-10.

Components

Package	Items	Quantity
	Bst DNA Polymerase (Large Fragment) (8 U/μL)	1 vial (1,600 U)
1,600 U	10× Bst DNA Polymerase Reaction Buffer	1 vial (1 mL)
	100 mM MgSO ₄	1 vial (0.4 mL)
	Bst DNA Polymerase (Large Fragment) (8 U/μL)	1 vial (8,000 U)
8,000 U	10× Bst DNA Polymerase Reaction Buffer	1 vial (1 mL)
	100 mM MgSO₄	1 vial (0.4 mL)



Specifications

Expression system

Escherichia coli

Concentration

8 U/µL

Purity

>98% as determined by SDS-PAGE analysis.

Form

Liquid

Application

Loop-mediated Isothermal Amplification (LAMP)

Buffer

Bst DNA Polymerase (Large fragment) is supplied in 10 mM Tris-HCl (pH7.5), 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1% Triton® X-100 and 50% Glycerol.

Unit Definition

One unit is defined as the amount of the enzyme incorporates 10 nmol of dNTP into acidinsoluble product in 30 minutes at 65°C.

Instruction

Shipping

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

Stability & Storage

This product is stable after storage at:

 -20°C or -80°C for 12 months under sterile conditions from date of receipt.

Avoid repeated freeze/thaw cycles.

Disclaimer: For Research Use or Further Manufacturing Only.